




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C12N 15/38, 7/04, A61K 39/245, 39/27	A1	(11) International Publication Number: WO 00/08165 (43) International Publication Date: 17 February 2000 (17.02.00)
(21) International Application Number: PCT/EP99/05476 (22) International Filing Date: 26 July 1999 (26.07.99) (30) Priority Data: 98202594.2 31 July 1998 (31.07.98) NL (71) Applicant (for all designated States except US): AKZO NOBEL N.V. [NL/NL]; Velperweg 76, NL-6824 BM Arnhem (NL). (72) Inventors; and (75) Inventors/Applicants (for US only): SONDERMEIJER, Paulus, Jacobus, Antonius [NL/NL]; Mahonie 21, NL-5831 RN Boxmeer (NL). VISSER, Nicolaas [NL/NL]; De Sering 26, NL-5831 RV Boxmeer (NL). DHORE, Cherida, Rachel [NL/NL]; Aert van der Neerstraat 5, NL-5831 VR Boxmeer (NL). (74) Agent: OGILVIE-EMANUELSON C., M.; P.O. Box 20, NL-5340 BH Oss (NL).		(81) Designated States: AL, AU, BA, BB, BG, BR, CA, CN, CU, CZ, EE, GE, HU, ID, IL, IN, IS, JP, KP, KR, LC, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, SL, TR, TT, UA, US, UZ, VN, YU, ZA, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>

(54) Title: ATTENUATED EQUINE HERPESVIRUS**(57) Abstract**

The present invention relates to novel Equine herpesvirus (EHV) mutants comprising one or more deletions, substitutions, or insertions in the endogenous promoter region of an essential viral gene, preferably the immediate early gene of EHV. The EHV mutants are stable and have reduced virulence, which makes them very suitable for use in a live vaccine. The invention furthermore relates to live vaccines comprising said EHV mutants, to DNA sequences and vectors harbouring a mutated EHV sequence, to host cells transfected with said DNA or vectors. The invention also relates to a method of attenuating EHV in general, and EHV-1 in particular.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

ATTENUATED EQUINE HERPESVIRUS

The present invention relates to a new Equine herpesvirus (EHV) mutant, to its use in vaccines, to DNA sequences and vectors harbouring a mutated EHV sequence, to host cells transfected with such vectors and to a method of attenuating EHV in general, and EHV-1 in particular.

Equine herpesviruses comprise a group of antigenic distinct biological agents which cause a variety of infections in horses, ranging from subclinical to fatal disease.

The Equine herpesvirus is inter alia one of the most common causes of infectious abortions and may account for 15% of all mare abortions that occur during the last six months of pregnancy. Equine herpesvirus type 1 (EHV-1) can cause abortion, perinatal foal mortality, respiratory inflammation and neurologic disease.

Although very similar to EHV-4 (previously classified as EHV-1 subtype 2), EHV-1's main clinical sign is abortion while EHV-4 results in respiratory disease. But both cross over and can be clinically indistinguishable as a disease.

The virus is contracted through inhalation. The respiratory signs can range from severe to inapparent. Abortion occurs primarily in mares over 7 months pregnant, follows infection in 14 to 120 days and can occur suddenly without signs in the mare. The virus infects fetal lung and liver tissue and the mare placental endothelial tissue. Abortion may occur because of direct effects on the fetus or because of placental separation. Near term fetuses may be born alive but die quickly due to lung pathology.

Primary infection of upper respiratory tract of young horses results in a febrile illness which lasts for 8 to 10 days. Immunologically experienced mares may be reinfected via the respiratory tract without disease becoming apparent so that abortion usually occurs without warning. The neurological syndrome is associated with respiratory disease or abortion and can affect animals of either sex at any age, leading to incoordination, weakness and posterior paralyses.

Other EHV viruses are EHV-2 or Equine Cytomegalovirus, which is a ubiquitous antigenically heterogeneous, usually slowly growing group of viruses, causing no-known disease, and EHV-3, the Equine Coital exanthema virus which is the causative agent of a relatively mild pregenital exanthema of both mare and stallion. EHV-1 subtype 2 is now called EHV-4, and is primarily associated with respiratory disease, although sporadic EHV-4 induced abortions have been reported.

The genomic structure of the EHV is similar to that of other alpha herpesviruses, comprising a double stranded linear molecule consisting of two covalently linked segments (U_L and U_S), wherein the U_S segment is flanked by inverted repeats. Of the EHV EHV-1 is the most extensively studied. Telford, E.A.R. et al., Virology 189, 304-316 (1992) have published the complete DNA sequence of EHV-1. The genome consists of about 150,000 bp and about 76 distinct genes have been recognised up to now.

Equine herpesviruses in general and EHV-1 in particular, are ubiquitous pathogens in horses. EHV-1 can even cause epidemics of abortion, respiratory tract disease and central nervous system disorders. Prevention of infection with the virus is therefore of major economic importance, because EHV can be a severe threat, especially to horses living in close groups, such as studs.

Current vaccines against these viruses comprise chemically inactivated viruses or attenuated live viruses. However, these require multiple administration and have only a limited efficacy.

Inactivated vaccines general induce only a low level of immunity. Attenuated live vaccines are thus preferred because they evoke a more long-lasting immune response and are easier to produce. Attenuation can be obtained by serial passages of virulent strains in tissue culture of other hosts than the natural one. However, the strains thus obtained are not very well defined and were considered not to be effected. Furthermore, there is always a risk that the viruses revert to virulence resulting in disease in inoculated animals. For this reason genetic attenuation was adopted as a novel approach to obtain safe vaccine strains.

Genetic attenuation consists for example of deletion of one or more non-essential genes. Examples are viruses with deletions in the thymidine kinase gene or the gene encoding glycoprotein gE. Both have been used successfully for genetic attenuation of herpesviruses in general but similar mutations in EHV-1 strongly abolished replication in the host and made the viruses of no longer use for vaccine purposes. The mutations furthermore often did not result in the desired level of attenuation.

It is therefore a first objective of the present invention to provide new EHV virus strains that do not have the drawbacks mentioned above when used in vaccines.

It is a second objective of the invention to provide a method for the more general genetic attenuation of herpesviruses.

The third objective of the invention is to provide for live vaccines that comprise said novel EHV mutant viruses.

The first objective of the present invention is achieved by an Equine herpesvirus (EHV) mutant, comprising one or more deletions, substitutions or insertions in the
5 endogenous promoter region of an essential viral gene. Preferably said mutation is a deletion in said endogenous promoter region.

Because attenuation of EHV by mutating non-essential genes was not successful, at least for vaccine purposes, the skilled person would have considered a mutation in the endogenous promoter region of essential genes to be even less promising since one would
10 not have expected the virus to replicate in the host at all. Therefore, the finding of the invention that the endogenous promoter region of essential genes can be mutated to reduce the virulence of EHV viruses leading to a successful attenuation is highly surprising.

The mutations according to the invention do not abolish replication of the virus in the host and affects the level of virulence of the virus. The EHV mutants according to the
15 invention are stable and attenuated with respect to the parent strain, which make them very suitable for use in a live vaccine.

In a first embodiment of the invention the promoter region is of the immediate early (IE) gene of the EHV virus. Other essential genes of which the promoters can be used are the genes involved in DNA replication (e.g. gene 57 in EHV-1 encoding the helicase/primase
20 complex), or transactivators of immediate early genes such as gene 12 in EHV-1 or genes encoding essential structural components of the virion such as the glycoproteins gB, gH and gL in EHV-1 or other herpesviruses.

The sequence upstream of the transcription initiation start contains several elements that regulate expression of a gene. All these elements together are defined as the promoter.
25 The core sequence of the promoter also known as TATA-box is located at about 40 base pairs from the transcription start although this structure may vary considerably from case to case. Other more or less conserved elements are located further upstream such as the CCAAT sequence and CG repeated motives. Some elements are specifically found in certain types of promoters such as a TAATGARATTC consensus sequence in immediate early
30 genes of herpesviruses. Also the sequence corresponding to the non-translated region upstream of the ATG start codon can be involved in promoter activity. Many of these

elements have not yet been defined nor is fully understood how they interact with the so-called transcription factors as a result of which transcription of the gene is regulated.

For the purpose of the present invention however, a promoter is defined as the sequence extending over several hundreds of base pairs upstream of the coding region of the gene with the most important part situated in the 200 base pairs preceding the transcription initiation site. Examples of sequences that contain promoters from essential genes can be retrieved from GenBank. The complete sequence for EHV-1 and EHV-4 have been deposited at GenBank under Accession No. M86664 and AF030027, respectively. For genes 57 and 12 from EHV-1, the promoter will be located between nucleotides 101.600-102.347 and 12.900-13.505, respectively.

Alternatively, promoter regions can be determined by nucleotide sequence analysis of DNA fragments that contain an essential gene including the region upstream of the coding sequence.

A preferred embodiment of the invention is the endogenous promoter of the IE gene of EHV-1 of which one copy is located between nucleotide 118.590 and about 119.890 in the Inverted Repeat of the short segment (IRs).

It has been found according to the invention that the endogenous promoter region of essential genes can be more suitable for attenuating the virulence of EHV. Until now this has not been considered because the manipulation of an essential gene was assumed to be deleterious for efficient replication in the host.

The term "essential gene" as used herein is intended to encompass genes that are obligatory for replication in the host. "Gene" is used solely for the coding sequence, whereas "promoter region" refers to the regulatory sequences necessary for expression of the gene.

Preferred mutations according to the invention are deletions between 1 and about 500 bases at any position within the promoter region that reduce the level of expression of the essential gene that is located downstream and attenuates virulence of the pathogen. By no means does a deletion according to the present invention abolish the expression of the gene product.

For EHV-1 preferred deletions are deletion of the SacI-SacI fragment or the HindIII-ClaI fragment or the NdeI-NdeI fragment or the SphI-SphI fragment of the promoter region of the Immediate Early gene. Highly preferred is a deletion of the HindIII-ClaI fragment of the IE promoter region; this deletion abolishes virulence of the virus without effecting local

replication in the host. An EHV-1 mutant comprising a deletion of the HindIII-ClaI fragment of the promoter region of its IE gene shows exceptionally good level of attenuation without any negative effect on its replication, thus making this mutant highly suitable for use in a live vaccine.

5 As an alternative fragments with a size between 1 and about 500 bases can be inserted in the endogenous promoter region, provided that said insertion does not abolish expression of the essential gene product. Such insertions according to the invention do not interfere with the virus ability of replication and have the advantage that they can lead to a desired level of reduction in virulence.

10 Substitutions can be made in the endogenous promoter of an essential viral gene varying from one or more nucleotides up to 500 base pairs provided that said substitution does not lead to loss of expression of the essential gene product. Substitutions according to the invention do not result in abolished replication of the virus in the host, but can lead to a desired level of attenuation.

15 Substitution according to the invention should not be confused with substitution of the complete promoter region by a cell, tissue or host specific promoter as described by Glazenburg et al. in US-5,580,564. Glazenburg aims at choosing the cell, the tissue and/or the host such that the harmful characteristics of the micro-organism are not expressed or are only expressed to an acceptable degree. The overall virulence of the virus is not decreased,
20 but instead the localisation of the viral replication is limited to particular cells, tissues or hosts. Substitutions according to the invention are not intended to encompass such mutations into specific promoters that lead to a modified cell, tissue or host tropism.

Deletions can be introduced by the following methods. Based on the map of the promoter region, suitable restriction sites can be deduced to remove fragments of a defined
25 size and at a defined position in the promoter. As an alternative one could start at a single site, and make progressive deletions in either one or two directions using the technique described by Henikoff (Gene 28, 351-359 (1984)). Also linker or PCR-mediated mutagenesis (Current Protocols in Molecular Biology, Eds. Ausubel et al., Chapter 8, John Wiley & Sons, Inc. (1996) can be used.

30 Mutations introduced in cloned subfragments can be transferred to the virus genome as for example described in Maniatis, T. et al. (1982) in "Molecular cloning", Cold

Spring Harbor Laboratory; European Patent Application 74.808; Roizman, B. & Jenkins, F.J. (1987), Science 229, 1208; Higuchi, R. et al. (1988), Nucleic Acids Res. 16, 7351.

Briefly, this can be accomplished by constructing a recombinant DNA molecule for recombination with Equine herpesvirus DNA. Such a recombinant DNA molecule comprises
5 vector DNA which may be derived from any suitable plasmid, cosmid, virus or phage, and contains Equine herpesvirus DNA of the region identified above.

Examples of suitable cloning vectors are plasmid vectors such as pBR322, the various pUC and Bluescript plasmids, cosmid vectors, e.g. THV, pJB8, MUA-3 and CosI, bacteriophages, e.g. lambda-gt-WES-lambda B, charon 28 and the M13mp phages or viral
10 vectors such as SV40, Bovine papillomavirus, Polyoma and Adeno viruses. Other vectors to be used can be retrieved from the Intelligenetics vector database accessible through website <http://www.seqnet.dl.ac.uk>.

A deletion to be introduced in the described region can be incorporated first in a recombinant DNA molecule carrying the promoter region of the essential gene of EHV by
15 means of a restriction enzyme digest with one or more enzymes of which the cleavage sites are correctly positioned in the promoter region of the gene. Recircularization of the remaining recombinant DNA molecule would result in a derivative lacking at least part of the promoter region. Alternatively, progressive deletions can be introduced either in one or two directions starting from within the restriction enzyme cleavage site present within the sequence of the
20 gene. Enzymes such as Bal31 or exonuclease III can be used for this purpose.

Recircularized molecules are transformed into E. coli cells and individual colonies can be analyzed by restriction mapping in order to determine the size of the deletion introduced into the promoter region. An accurate positioning of the deletion can be obtained by sequence analysis.

25 In case the insertion of a heterologous nucleic acid sequence is desired the recombinant DNA molecule comprising the EHV essential gene may be digested with appropriate restriction enzymes to produce linear molecules whereafter the heterologous nucleic acid sequence can be ligated to the linear molecules followed by recircularization of the recombinant DNA molecule.

30 Optionally, a deletion is introduced into the promoter region of the EHV gene concomitantly with the insertion of the heterologous nucleic acid sequence.

In case the method of in vivo homologous recombination is applied for the preparation of an EHV mutant according to the invention the EHV sequences which flank the deleted gene sequences or the inserted heterologous nucleic acid sequences should be of appropriate length, e.g. 50-3000 bp, as to allow in vivo homologous recombination with the viral EHV genome to occur.

Subsequently, cells, for example equine cells such as equine dermal cells (NBL-6) or cells from other species such as RK13, Vero and BHK cells can be transfected with EHV DNA in the presence of the recombinant DNA molecule containing the mutation flanked by appropriate EHV sequences whereby recombination occurs between the EHV sequences in the recombinant DNA molecule and the corresponding sequences in the EHV genome.

Recombinant viral progeny is thereafter produced in cell culture and can be selected for example genotypically or phenotypically, e.g. by hybridization. The selected EHV mutant can be cultured on a large scale in cell culture whereafter EHV mutant containing material can be collected therefrom.

The EHV-1 genome contains two copies of the Immediate Early gene, in opposite orientation. In this case a convenient method for mutating the promoter is the construction of a set of cosmids harbouring fragments of the EHV genome, including only one of the two copies of the IE gene and its promoter region. This single copy of the promoter region can then be mutated in a routine manner. The set of cosmids is transfected to a confluent monolayer of host cells. Viruses will be formed by recombination and can be recovered from plaques formed in the monolayer. It is necessary to use one copy of the IE promoter region and corresponding gene for the preparation of a mutant according to the invention to prevent replacement of the mutation by the non-mutated promoter region. The mutant virus will restore a second identical copy of the mutated promoter region of the IE gene while assembling a functional full-size genome containing both inverted repeats flanking the Us.

It is standard procedure to test the efficacy and safety of a vaccine in vivo. Initial testing of mutant EHV viruses can be performed in the well-known mouse model as described by van Woensel et al., J. Virological Methods 54, 39-49, (1995) and Osterrieder et al., Virology 226, 243-251, (1996). Final testing of the safety and efficacy of the virus strain takes place in vivo in horses.

The mutation(s) in the endogenous promoter of one or more essential genes can be combined with one or more mutations in one or more other genes and/or their promoters.

Examples of these are mutations creating a marker vaccine which allow differentiation of vaccinated animals from infected animals. This way a further optimisation of the vaccine's properties can be obtained.

5 The invention further relates to a nucleic acid sequence, comprising the endogenous promoter region of an EHV essential gene and optionally one or more flanking sequences, which promoter region comprises one or more deletions, substitutions or insertions. The gene is for example an Immediate Early gene, or any other essential gene, such as those described hereinabove.

10 Specific deletions are deletions of the SacI-SacI fragment or the HindIII-ClaI fragment or the NdeI-NdeI fragment or the SphI-SphI fragment of the promoter region of the Immediate Early gene. Particularly preferred are deletions of the HindIII-ClaI fragment of the promoter region of the Immediate Early gene of EHV.

The nucleic acid sequence in a specific embodiment comprises the promoter region of EHV-1 or EHV-4, more specifically EHV-1.

15 This nucleic acid sequence can be incorporated in a recombinant DNA molecule, which is also part of the invention. The selection of suitable vectors to prepare the recombinant DNA molecule are well within the reach of the person skilled in the art.

In addition the invention relates to a host cell harbouring such a recombinant DNA molecule.

20 The present invention furthermore provides for a vaccine comprising an EHV mutant of the invention and a pharmaceutically acceptable carrier or diluent.

A live attenuated mutant according to the present invention can be used to vaccinate equines, particularly domestic and non-domestic, and more specifically horses. Vaccination with such a live vaccine is preferably followed by replication of the mutant within
25 the inoculated host, which host will then elicit an immune response against EHV and the animal inoculated with the EHV mutant according to the invention will be immune to infection by EHV. Thus, a mutant according to claim 1 can serve as a live vaccine.

For the preparation of a live vaccine the recombinant EHV mutant according to the present invention can be grown on a cell culture, for example of equine, rabbit, hamster or
30 calf origin. The viruses thus grown can be harvested by collecting the tissue cell culture fluids

and/or cells. The live vaccine may be prepared in the form of a suspension or may be lyophilised.

The vaccine according to the invention can be prepared using standard techniques available in the art. In general the vaccine is prepared by mixing the virus with a
5 pharmaceutically acceptable carrier or diluent.

For administration to animals, the EHV mutant according to the present invention can be given inter alia intranasally, intradermally, subcutaneously or intramuscularly.

Pharmaceutically acceptable carriers or diluents that can be used to formulate a vaccine according to the invention are sterile and physiologically compatible such as for
10 example sterile water, saline, aqueous buffers such as alkali metal phosphates (e.g. PBS), alcohols, polyols and the like. In addition the vaccine according to the invention may comprise other additives such as adjuvants, stabilizers, anti-oxidants, preservatives and the like.

Suitable adjuvants include but are not limited to aluminium salts or gels, carbomers,
15 non-ionic blockcopolymers, tocopherols, monophospheryl lipid A, muramyl dipeptide, oil emulsions (w/o or o/w), cytokines. The amount of adjuvant added depends on the nature of the adjuvant itself.

Suitable stabilizers for use in a vaccine according to the invention are for example carbohydrates including sorbitol, mannitol, starch, sucrose, dextrin and glucose, proteins
20 such as albumin or casein, and buffers like alkaline phosphates.

Suitable preservatives include, amongst others, thimerosal, merthiolate and gentamicin.

Live vaccines according to the invention comprise an effective amount of the aforementioned EHV mutant virus and a pharmaceutically acceptable carrier. The term "effective"
25 as used herein is defined as the amount sufficient to induce an immune response in the target animal. The amount of virus will depend on the route of administration, the time of administration as well as age, general health and diet of the subject to be vaccinated.

The dosages in which the live vaccines according to the invention can prevent infectious disease can be readily determined by routine trials with appropriate controls and
30 are well within the routine skills of the practitioner.

The useful dosage to be administered will vary depending on the age, weight, mode of administration and type of pathogen against which vaccination is sought. A suitable dosage can be for example about $10^{3.0}$ - $10^{7.0}$ pfu/animal.

According to a further aspect thereof the invention provides a process for the preparation of EHV mutants, comprising transfecting a cell culture with a recombinant DNA molecule and EHV genomic DNA.

The invention thus provides for a method of genetically attenuating equine herpesviruses, comprising mutation of the endogenous promoter region of an essential gene, which mutation consists of one or more deletions, substitutions or insertions in the promoter region of an essential gene.

The present invention will be further illustrated in the non-limiting examples that follow. All molecular biological techniques that are used in the examples are standard procedures as for example described in Sambrook et al., "Molecular Cloning, A Laboratory Manual", Cold Spring Harbor Laboratory Press (1989). These techniques will not be described in detail.

LEGENDS

Figure 1 shows a map of the EHV-1 genome with the position of five cosmid DNA inserts that restore a viable virus after cotransfection and recombination between their overlapping sequences. Deletions in the IE gene promoter were introduced in a 8 kb SpeI fragment of the IRs that was re-assembled with a 21 kb SpeI and used as such to replace cosmid 2D9 in the cotransfection experiment.

Figure 2 shows a schematic layout of the promoter region of the IE gene in the IRs from EHV-1. Various sequence elements that may regulate transcription initiation are indicated. Below are shown the deletions that were introduced and the mutants that were generated by incorporating these deletions into the viral genome after co-transfection with the set of overlapping cosmid DNA inserts. Strain EHV24 was restored in a similar manner but contained no deletions in the promoter region.

EXAMPLES**EXAMPLE 1: Construction of the cosmid set for generating EHV-1 virus**

The SuperCos 1 cosmid vector kit was purchased from Stratagene (Catalog# 251301) and enzymes were purchased at New England Biolabs. The vector was further
5 modified by adding extra restriction enzyme sites to it. For this a DNA linker was purchased from Pharmacia containing the following restriction sites: BamHI, I-SceI, PacI, Ascl, EcoRV, PacI, Ascl, I-SceI and BamHI. The SuperCos 1 vector and the linkers were both cut with BamHI. The BamHI digested vector was dephosphorylated with alkaline phosphatase after which the BamHI digested linker was ligated into the SuperCos 1 vector by T4 DNA ligase,
10 all according to the manufacturer's instructions. The resulting vector was designated THM and used for cloning the EHV-1 inserts.

Viral DNA was obtained from the EHV-1 M8 strain, a pathogenic EHV-1 strain isolated from a horse with severe signs of an EHV-1 infection, but could be any pathogenic EHV-1 strain. This particular virus was grown in cultures of Vero cells and was recovered by
15 centrifugation. The viral DNA was prepared by lysis with EDTA and SDS, phenol extraction and alcohol precipitation.

The first set of cosmids was constructed by digestion of the EHV-1 DNA with PacI. After phenol extraction of the M8 PacI digests, the ends were filled in with T4 DNA polymerase and then dephosphorylated with alkaline phosphatase. The cosmid vector THM
20 was digested with EcoRV and the inserts were ligated into the vector with T4 DNA ligase.

The ligation mix was added to a packaging mix (Gigapack, Stratagene) according to the manufacturers instructions. The packaged DNA was added to a fresh overnight culture of E.coli DH1 and placed for 1 hour at 37°C. The bacteria suspension was then spread onto agar plates containing ampicillin. Plasmid DNA from all colonies was submitted to restriction
25 enzyme analysis on the insert.

For the construction of other cosmids the same procedure was followed only now the viral DNA was digested with Ascl, Asel, Rsrl, or NotI, all ends were then filled in with T4 DNA polymerase and the inserts ligated into the EcoRV site of the vector.

To obtain a third generation of cosmids the viral DNA was sheared twice through a
30 19G needle, the ends were then filled in with T4 DNA polymerase and after phenol extraction and precipitation the inserts were cloned again into the EcoRV site of the cosmid vector

THM. The vector with the inserts was then packed, put on bacteria and the colonies analyzed.

From all colonies obtained the restriction maps were determined by multiple digestions. Then the location of the different clones was determined by comparing the restriction map of the clones with the restriction map of EHV-1 strain Ab4 (Telford et al, 1992).

Cosmids 2D3, 1A12, 1F4, 2C12 and 2D9 were selected such that the complete genome could be reconstituted by recombination between overlapping sequences (Fig. 1).

Viable virus was generated after transfecting EHV-1 inserts excised from the cosmids by digestion with I-SceI (Boehringer) into cultures of equine dermal cells (ATCC number CCL-57) using the calcium phosphate method. Viral progeny was passaged and plaque purified on monolayers of rabbit kidney (RK13, ATCC number CCL-37) or other kidney cells that support viral replication.

EXAMPLE 2: *Introducing deletions in the promoter of the IE gene*

The genome of EHV-1 contains two copies of the Immediate Early (IE) gene located in the repeated sequence elements IRs and TRs flanking the unique short region. Stable mutations affecting expression of the IE gene need to be introduced at both positions simultaneously before the new phenotype will be expressed.

Therefore, one of the cosmids named "2D9" was modified such that a fragment of about 7 kb at the right end of the TRs and containing a single copy of the IE gene, was deleted. This was obtained by subcloning the two SpeI fragments of 8 and 21 kb from 2D9 in pGEM-9Zf(-) (Promega) using the SpeI or SpeI/XbaI sites, respectively (Fig. 1). Insertion of the 21 kb fragment resulted in pEHV17 and restored a single SpeI site in which the 8 kb SpeI fragment subsequently can be inserted to regenerate a 29 kb EHV-1 fragment used for replacing 2D9 in the cosmid transfection described in Example 1. The 8 kb SpeI contains all of the coding sequence of the IE gene and about 2 kb of upstream sequences including the promoter of the IE gene. Mutations in the promoter region were introduced in a plasmid designated pEHV06 that contained a 2 kb subfragment from the 8 kb SpeI. This insert included a 1 kb EcoRV fragment which essentially represented the promoter and other regulatory elements located in the sequence upstream of the coding region of the IE gene (Fig. 2).

By selecting the proper restriction enzymes that had two site in the EcoRV fragment of pEHV06 and relatively near to each other, small deletions with an exact size could be introduced at various positions in the promoter sequence of the IE gene. The enzymes SacI, NdeI, SphI and a combination of ClaI and HindIII were suitable for this purpose. The deletions introduced in pEHV06 by digesting with each of these enzymes and recircularizing the plasmid DNA with T4 DNA ligase are represented in Fig. 2.

The mutated EcoRV inserts were exchanged with the original EcoRV insert in the 8 kb SpeI fragment and these were subsequently assembled with the 21 kb SpeI in pEHV17 resulting in plasmids that were designated pEHV22, pEHV23, pEHV25 and pEHV26 for the deletions generated with the enzymes SacI, HindIII/ClaI, NdeI and SphI, respectively (Fig. 2). pEHV24 contained the re-assembled 8 and 21 kb SpeI fragments but without any deletion in the promoter region. These plasmids were used to replace 2D9 in the cotransfection of equine dermal cells with the cosmid inserts from 2D3, 1A12, 1F4 and 2C12, generating viable virus with defined small deletions in the promoter region of both copies of the IE gene. Viral progeny was plaque purified and amplified on RK13 cells. Presence and size of the deletions were confirmed by DNA blot analysis and PCR.

Mutant strains were designated EHV22, EHV23, EHV25 or EHV26 with the number corresponding to the plasmid used in the cotransfection experiment. Strain EHV24 represents the control virus and contains the non-mutated promoter region.

Cosmid derived mutant EHV strains 22, 25 and 26 and the control strain EHV24 grow to titers between 2 and 4×10^6 pfu/ml on RK13 cells, which is slightly less than the parent M8 strain that grows to titer of 5×10^6 pfu/ml. Strain EHV23 replicated less efficiently and resulted in titers of about 2×10^5 pfu/ml.

EXAMPLE 3: Analysis of virulence in mice

Infection with EHV-1 can cause pyrexia and clinical signs of respiratory disease but most important is the induction of abortion. However, seen the complexity of vaccination trials in pregnant horses, models have been developed in mice measuring the pathogenicity of EHV-1 strains based on weight reduction (van Woensel et al., J. Virological Methods 54, 39-49, (1995); Osterrieder et al., Virology 226, 243-251, (1996).

Mice that are 4-5 weeks of age are inoculated intranasally with a dose of $10^{6.5}$ pfu per animal. Body weights are determined on the 9 days following infection and the weight

gain as percentage of the weight at day 0 are plotted in a graph. The degree of pathogenicity of various EHV-1 strains is deduced from the position of resulting curves relative to each other. Significant losses in body weight, particularly at day 3 and 4 after infection, correlate with high virulence of the strain of interest. Series of EHV-1 isolates carrying deletions in the IE promoter region (see Example 2) can be analyzed for virulence by comparison with the parent M8 strain or cosmid generated EHV-1 virus containing the intact promoter sequence.

EXAMPLE 4: Analysis of virulence in horses

Although pathogenicity of individual EHV-1 strains using the mouse model can be correlated with behavior in the natural host, more conclusive evidence can be obtained from vaccination trials in horses. In addition, the level of protection against challenge infection and ultimately in preventing abortion in pregnant mares can only be established in the target animal.

The experiment was performed in four conventional colts at the age of about one year and previously tested for the absence of EHV-1 serum antibodies. Animals were divided in groups of two, one group receiving mutant strain EHV23, the other receiving mutant strain EHV24. The colts were inoculated intranasally with a single dose of about 5×10^5 pfu, reconstituted in 2 ml Unisolve diluent and applying 1 ml in each nostril. Following 10 days, the animals were monitored for clinical reactions. Antibodies were measured at several time points either by virus-neutralisation or using a complement-fixing antibody assay. Infected blood leucocytes (viremia) and excreted virus in nasal swabs were determined by titration.

After three weeks, a fifth EHV-1 seronegative horse was added and two weeks later all five horses were challenged intranasally with 7×10^5 pfu of the pathogenic EHV-1 AB4p strain. The animals were again monitored for clinical reactions, virus excretion and infected blood leucocytes.

The horses vaccinated intranasally with a high dose of mutant strain EHV23 showed virtually no signs of disease, whereas horses vaccinated with mutant control strain EHV24 showed severe signs of EHV-1 infection including fever and ocular or nasal discharges during several days. Both mutant strains were obtained in an identical way, in case of EHV23 introducing a 160 bp deletion in the IE promoter and in case of EHV24 no deletion at all (see example 2). EHV24 strain thus represents the control virus, containing the non-mutated IE

promoter. Surprisingly, both strains were replicating in the nasal cavity, although it appeared that replication of the EHV23 strain was slightly delayed.

Challenge of all horses with EHV-1 strain AB4p did not induce significant signs of disease in the animals that had been previously inoculated with strain EHV23 or EHV24. No
5 viremia was detected in any of the horses inoculated with either strain EHV23 or EHV24 which indicates a solid systemic protection of these animals against a challenge infection with the EHV-1 AB4p strain.

CLAIMS

1. Equine herpesvirus (EHV) mutant, comprising one or more deletions, substitutions or insertions in the endogenous promoter region of an essential viral gene.
2. EHV mutant as claimed in claim 1, wherein deletions are introduced into the promoter.
3. EHV mutant as claimed in claims 1-2, wherein the gene is the Immediate Early gene.
4. EHV mutant as claimed in claims 1-3, wherein the mutant virus is the EHV-1 virus or the EHV-4 virus.
5. EHV mutant as claimed in claims 1-4, further comprising one or more mutations in one or more other genes and/or their promoters.
6. EHV-1 mutant as claimed in claims 1-5, comprising a deletion of the SacI-SacI fragment or the HindIII-ClaI fragment or the NdeI-NdeI fragment or the SphI-SphI fragment of the promoter region of the Immediate Early gene.
7. Nucleic acid sequence, comprising the endogenous promoter region of an essential gene from EHV and optionally one or more flanking sequences, which promoter region comprises one or more deletions, substitutions or insertions.
8. Nucleic acid sequence as claimed in claims 7, wherein the gene is an Immediate Early gene.
9. Nucleic acid sequence as claimed in claim 8, comprising a deletion of the SacI-SacI fragment or the HindIII-ClaI fragment or the NdeI-NdeI fragment or the SphI-SphI fragment of the promoter region of the Immediate Early gene.
10. Nucleic acid sequence as claimed in claims 7-9, wherein the EHV is EHV-1 or EHV-4.
11. Recombinant DNA molecule comprising a nucleic acid sequence as claimed in claims 7-10.
12. Host cell harbouring a recombinant DNA molecule as claimed in claim 11.
13. Vaccine comprising an EHV mutant as claimed in claims 1-6 and a pharmaceutically acceptable carrier or diluent.
14. A process for the preparation of an EHV mutant as claimed in claims 1-6, comprising transfecting a cell culture with a recombinant DNA molecule as claimed in claim 11 and EHV genomic DNA.

15. Method of genetically attenuating EHV, comprising mutation of the endogenous promoter region of an essential gene, which mutation consists of one or more deletions, substitutions or insertions in the promoter region of an essential gene.
16. Method as claimed in claim 15, wherein the EHV is EHV-1 or EHV-4.
- 5 17. Method as claimed in claims 16-17, wherein the gene is an Immediate Early gene.




FIG.1




FIG.2

INTERNATIONAL SEARCH REPORT

Internat I Application No

PCT/EP 99/05476

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/38 C12N7/04 A61K39/245 A61K39/27

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N A61K C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	LEWIS JB ET AL: "Transcriptional control of the equine herpesvirus 1 immediate early gene" VIROLOGY, vol. 197, no. 2, December 1993 (1993-12), pages 788-792, XP002125016 ORLANDO US Plasmid pIEbetagal containing the EHV-1 Immediately Early promoter. page 788, right-hand column, last paragraph	7,8
A	WO 92 01045 A (THE UNIVERSITY OF GLASGOW) 23 January 1992 (1992-01-23) example 3 --- -/--	1-8, 10-12

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

8 December 1999

Date of mailing of the international search report

18/01/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Cupido, M

INTERNATIONAL SEARCH REPORT

Intern: al Application No

PCT/EP 99/05476

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>SMITH R H ET AL.: "Nuclear localization and transcriptional activation activities of truncated versions of the Immediate-Early gene product of Equine Herpesvirus 1"</p> <p>JOURNAL OF VIROLOGY, vol. 69, no. 6, June 1995 (1995-06), pages 3857-3862, XP002088567</p> <p>the whole document</p> <p>----</p>	1-17
A	<p>MARSHALL K R ET AL: "AN EQUINE HERPESVIRUS-1 GENE 71 DELETANT IS ATTENUATED AND ELICITS A PROTECTIVE IMMUNE RESPONSE IN MICE"</p> <p>VIROLOGY,US,ACADEMIC PRESS,ORLANDO, vol. 231, no. 1, page 20-27 XP002055348</p> <p>ISSN: 0042-6822</p> <p>EHV-1 gene 71 deletion mutant is attenuated and may be used as vaccine strain (see also W098/26049)</p> <p>the whole document</p> <p>----</p>	1-17
A	<p>W0 96 04394 A (MEDICAL RESEARCH COUNCIL)</p> <p>15 February 1996 (1996-02-15)</p> <p>page 5, line 15 -page 6, line 25</p> <p>-----</p>	15-17

INTERNATIONAL SEARCH REPORT

Information on patent family members

Internal Application No

PCT/EP 99/05476

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
W0 9201045	A	23-01-1992	AU 8212891 A	04-02-1992
			CA 2086740 A	07-01-1992
			EP 0538299 A	28-04-1993
			HU 67778 A	28-04-1995
			NZ 238834 A	26-05-1992
W0 9604394	A	15-02-1996	AU 695405 B	13-08-1998
			AU 3119595 A	04-03-1996
			CA 2195965 A	15-02-1996
			EP 0804602 A	05-11-1997
			JP 10503372 T	31-03-1998